

Publications: 2011 - 2015

2011

Agbon CA, Akiyemi CO, Onabanjo OO, Okeke EC. Nutrient composition and phytate-zinc molar ratio of prepared foods consumed by rural preschool children. *Nigerian J Nutr Sciences* 2010;31:58-63.

Agbon CA, Ngozi EO, Onabanjo OO. Production and nutrient composition of fufu made from a mixture of cassava and cowpea flours. *J Culinary Science & Techn* 2010;8:147-57.

Agbon CA, Okeke EC, Akinyemi CO. Anaemia, nutritional status and parasitic infection among preschool children living in rural Nigerian villages. *Nigerian J Nutr Sciences* 2010;31:48-52.

Agbon CA, Okeke EC, Omotayo A. Nutrient intake of rural preschool children in southwest Nigeria. *Infant, Child and Adolescent Nutr* 2010;2:336-9.

Dastgiri S, Sharafkhani R, Gharaaghaji R, Ghavamzadeh S. Prevalence, influencing factors and control of food insecurity: a model in the northwest of Iran. *Asia Pac J Clin Nutr* 2011;20:613-7.

Dossa RAM; Ahouandjinou ERT; Houngbe FE. Evaluation of the suitability and acceptability of a newly designed infant flour for infant feeding in the district of Bopa in south of Benin. *African J of Food, Nutrition and Development (AJFAND)*, 2011;11 No.4.

Li L, Hu Y, Zhao X, Wang J, Muzhingi T, Suter PM, Yin SA, Tang G. Spirulina can increase the total body vitamin A storage of Chinese children. *Experimental Biology* 2010, Abstract 2010.

Long KZ, Garcia C, Ko G, Santos JI, Al Mamun A, Rosado JL, Dupont HL, Nathakumar N. Vitamin A modifies the intestinal chemokine and cytokine responses to norovirus infection in Mexican children. *J Nutr* 2011;141:957-63.

Long KZ, Santos JI, Rosado JL, Estrada-Garcia T, Haas M, Al Mamun A, DuPont HL, Nanthakumar NN. Vitamin A supplementation modifies the association between mucosal innate and adaptive immune responses and resolution of enteric pathogen infections. *Am J Clin Nutr* 2011;93:578-85.

Nguyen TH, Tang HK, Kelly P, van der Ploeg HP, Dibley MJ. Association between physical activity and metabolic syndrome: a cross sectional survey in adolescents in Ho Chi Minh City, Vietnam. *BMC Public Health* 2010;10.

Sranacharoenpong K, Hanning RM. Diabetes prevention education program for community health care workers in Thailand. *J Community Health* 2011; Epub ahead of print.

Sranacharoenpong K, Hanning RM. Developing a diabetes prevention education programme for community health-care workers in Thailand: formative findings. *Prim Health Care Res & Dev* 2011;12:357-69.

Tang KH, Nguyen HH, Dibley MJ, Sibbritt DW, Phan NT, Tran TM. Factors associated with adolescent overweight/obesity in Ho Chi Minh city. *Int J Pediatr Obes* 2010;5:396-403.

Wijaya-Erhardt M, Muslimatun S, Erhardt JG. Fermented soyabean and vitamin C-rich fruit: a possibility to circumvent the further decrease of iron status among iron-deficient pregnant women in Indonesia. *Public Health Nutr* 2011;24:1-12.

Yang H, Qian M, Ding W, Gebre-Medhin M, Yan Y, Chen Z. A comparison study of two

reference data for evaluation of thyroid function in pregnancy. *Tianjin Med J* 2011;39:299-302.

Yu B, Wang J, Suter PM, Russell RM, Grusak MA, Wang Y, Wang Z, Yin S, Tang G. Spirulina is an effective dietary source of zeaxanthin to humans. *Br J Nutr* 2011;12:1-10.

2012

Abizari AR, Moretti D, Schuth S, Zimmermann MB, Armar-Klemesu M, Brouwer ID. Phytic acid-to-iron molar ratio rather than polyphenol concentration determines iron bioavailability in whole-cowpea meal among young women. *J Nutr* 2012;142:1950-5.

Abizari AR, Moretti D, Zimmermann MB, Armar-Klemesu M, Brouwer ID. Whole cowpea meal fortified with NaFeEDTA reduces iron deficiency among Ghanaian school children in a malaria endemic area. *J Nutr* 2012;142:1836-42.

Htet M K, Dillon D, Akib A, Utomo B, Fahmida U, Thurnham D I. Microcytic anaemia predominates in adolescent school girls in the delta region of Myanmar. *Asia Pac J Clin Nutr* 2012;21:411-5.

Jiang H, Li M, Yang D, Wen LM, Hunter C, He G, Qian X. Awareness, intention, and needs regarding breastfeeding: findings from first-time mothers in Shanghai, China. *Breastfeeding Med* 2012; Epub ahead of print.

Ma Aiguo. Micronutrient status and effects of supplementation in anemic pregnant women in rural China. Published thesis Wageningen University, the Netherlands 2012; ISBN 978-94-6173-493-8:148 pp.

Macharia-Mutie CW, Van de Wiel AM, Moreno-Londono AM, Mwangi AM, Brouwer ID. Sensory acceptability and factors predicting the consumption of grain amaranth in Kenya. *Ecol Food Nutr* 2011;50:375-92.

Macharia-Mutie CW. Efficacy of amaranth grain (*Amaranthus cruentus*) on anaemia and iron deficiency in Kenyan pre-school children. Published thesis Wageningen University, the Netherlands 2012; ISBN 978-94-6173-291-0:168 pp.

Macharia-Mutie CW, Moretti D, Van den Briel N, Omusundi AM, Mwangi AM, Kok FJ, Zimmermann MB, Brouwer ID. Maize porridge enriched with a micronutrient powder containing low-dose iron as NaFeEDTA but not amaranth grain flour reduces anemia and iron deficiency in Kenyan preschool children. *J Nutr* 2012;142:1756-63.

Rahman MA, Rahman B, Ahmad MS, Blann A, Ahmed N. Blood and hair lead in children with different extents of iron deficiency in Karachi. *Environ Res* 2012;118:94-100.

Reinsma K, Bolima N, Fonteh F, Okwen P, Yota D, Montgomery S. Incorporating cultural beliefs in promoting exclusive breastfeeding. *African J of Midwifery & Women's Health* 2012;6:65-70.

Thakwalakwa C, Ashorn P, Phuka J, Cheung YB, Briend A, Puumalainen T, Maleta K. A lipid-based nutrient supplement but not corn-soy blend modestly increases weight gain among 6- to 18-month-old moderately underweight children in rural Malawi. *J Nutr* 2010;140:2008-13.

Trang NH, Hong TK, Dibley MJ. Cohort profile: Ho Chi Minh City Youth Cohort-changes in diet, physical activity, sedentary behaviour and relationship with overweight/obesity in adolescents. *BMJ Open* 2012;2:print 2012.

Trang NH, Hong TK, Dibley MJ. Active commuting to school among adolescents in Ho Chi Minh City, Vietnam: change and predictors in a longitudinal study, 2004 to 2009. *Am J*

Prev Med 2012;42:120-8.

Van Beusekom I, Vossenaar M, Montenegro-Bethancourt G, Doak C, Solomons NW. Estimation of exclusive breastfeeding rates among mother-infant dyads in Quetzaltenango, Guatemala varies by interview methods and time-frame. *Br J Nutr* 2012; to be published.

Vossenaar M, Garcia R, Solomons NW, Doak CM, Peters L, Campos Ponce M. Ritual fluids in relation to early child nutrition in Quetzaltenango, Guatemala. *Sight and Life* 2012;40-46.

Yang DL, Jiang H, Li M, Qian X. Intervention intention of using mobile phone text messages and the internet as an intervention tool to promote breastfeeding. *Chinese Journal of Maternal and Child Health* 2011;30:4704-6 (in Chinese).

2013

Abizari AR. Efficacy of iron fortified cowpea flour in improving iron status of schoolchildren in malaria endemic rural Ghana. Ph.D. Thesis, Wageningen University, Wageningen, NL, 2013:191pp.

Ayoya MA, Spiekermann-Brouwer GM, Traoré AK, Garza C. Effect on school attendance and performance of iron and multiple micronutrients as adjunct to drug treatment of *Schistosoma*-infected anemic schoolchildren. *Food and Nutrition Bulletin* 2012;33:235-41.

Egbi G, Steiner-Asiedu M, Saalia F, Ayi I, Ofosu W, Setorglo J. Iron stores and anaemia prevalence among school children fed cowpea-based food plus fishmeal with vitamin C-rich drink. *Ann Nutr Metab* 2013;63(suppl 1):1-1960 (PO1351).

Finkelstein JL, O'Brien KO, Abrams SA, Zavaleta N. Infant iron status affects iron absorption in Peruvian breastfed infants at 2 and 5 mo of age. *Am J Clin Nutr* 2013;98:1475-84.

Flax VL, Phuka J, Cheung YB, Ashorn U, Maleta K, Ashorn P. Feeding patterns and behaviors during home supplementation of underweight Malawian children with lipid-based nutrient supplements or corn-soy blend. *Appetite* 2010;54:504-11.

Hong TK, Trang NH, Dibley MJ. Prevalence of metabolic syndrome and factor analysis of cardiovascular risk clustering among adolescents in Ho Chi Minh City, Vietnam. *Prev Med* 2012;55:409-11.

Hong TK, Trang NH, van der Ploeg HP, Hardy LL, Dibley MJ. Validity and reliability of a physical activity questionnaire for Vietnamese adolescents. *Int J Behav Nutr Phys Act* 2012;9:1-7.

Htet MK, Fahmida U, Dillon D, Akib A, Utomo B, Thurnham DI. The influence of vitamin A status on iron-deficiency anaemia in anaemic adolescent schoolgirls in Myanmar. *Public Health Nutr* 2013 Oct 16:1-8. [Epub ahead of print]

Hu QZ, Jiang H, Li M, Wang H, Cheng J, Qiu J, Qu JQ, Gao YN, Qian X. Descriptive study on workplace support for breastfeeding in Shanghai (no electronic version available). *Chinese J of Maternal and Child Health Care* 2013;28:1778-81.

Jiang H, Li M, Wen LM, Hu QZ, Yang DL, He GS, Baur LA, Qian X. Effect of a short message service intervention on infant feeding practices in Shanghai, China: a prospective, community-based, controlled study. *Lancet* 2013;382:4.

Li L, Zhao X, Wang J, Muzhingi T, Suter PM, Tang G, Yin S. Spirulina can increase

total-body vitamin A stores of Chinese school-age children as determined by a paired isotope dilution technique. *JNS* 2012;1:1-7.

Ochola SA, Labadarios D, Nduati RW. Impact of counselling on exclusive breast-feeding practices in a poor urban setting in Kenya: a randomized controlled trial. *Public Health Nutr* 2013;16:1732-40.

Rahman MA, Rahman B, Ahmed N. High blood manganese in iron-deficient children in Karachi. *Public Health Nutr* 2013;16:1677-83.

Reinsma KR. The efficacy of an audio program and discussion guide in promoting exclusive breastfeeding in Cameroon, Africa. A dissertation in partial fulfillment of the requirements for the degree of doctor of public health in nutrition. Loma Linda University, Loma Linda, CA, USA 2012: 233pp.

Tang G, Suter PM. Vitamin A, nutrition, and health values of algae: spirulina, chlorella, and dunaliella. *J of Pharmacy & Nutr Sciences* 2011;1:111-8.

Tang G, Muzhingi T, Grusak M. Vitamin A equivalence of spirulina B-carotene in US men. *FASEB J* 2013;27:638.8.

Thakwalakwa CM, Ashorn P, Jawati M, Phuka JC, Cheung YB, Maleta KM. An effectiveness trial showed lipid-based nutrient supplementation but not corn-soya blend offered a modest benefit in weight gain among 6- to 18-month-old underweight children in rural Malawi. *Public Health Nutr* 2012;15:1755-62.

Thakwalakwa CM, Kuusipalo HM, Maleta KM, Phuka JC, Ashorn P, Cheung YB. The validity of a structured interactive 24-hour recall in estimating energy and nutrient intakes in 15-month-old rural Malawian children. *Matern Child Nutr* 2012;8:380-9.

Thakwalakwa CM, Ashorn P, Phuka JC, Cheung YB, Briend A, Maleta KM. Impact of lipid-based nutrient supplements and corn-soy blend on energy and nutrient intake among moderately underweight 8-18-month-old children participating in a clinical trial. *Matern Child Nutr* (in press).

Tofail F, Hamadani JD, Mehrin F, Ridout DA, Huda SN, Grantham-McGregor SM. Psychosocial stimulation benefits development in nonanemic children but not in anemic, iron-deficient children. *J Nutr* 2013;143:885-93.

Trang NH, Hong TK, Van der Ploeg HP, Hardy LL, Kelly PJ, Dibley MJ. Longitudinal physical activity changes in adolescents: Ho Chi Minh City Youth Cohort. *Med Sci Sports Exerc* 2012;44:1481-9.

Ziem N, Egbi G, Steiner-Asiedu M, Armar-Klemesu M. Prevalence and determinants of anemia among older school children in the Volta region of Ghana. *Ann Nutr Metab* 2013;63(suppl 1):1-1960 (PO1157).

2014

Diana R, Khomsan A, Sukandar D, Riyadi H. Nutrition extension and home garden intervention in Posyandu: Impact on nutrition knowledge, vegetable consumption and intake of vitamin A. *Pakistan J Nutr* 2014;13:88-92.

Egbi G, Steiner-Asiedu M, Kwesi FS, Ayi I, Ofosu W, Setorglo J, Klobodu SS, Armar-Klemesu M. Anaemia among school children older than five years in the Volta Region of Ghana. *Pan Afr Med J* 2014;17 (suppl 1):10.

Fahmida U, Kolopaking R, Santika O, Sriani S, Umar J, Htet MK, Ferguson E. Effectiveness on improving knowledge, practices, and intakes of "key problem nutrients" of a complementary feeding intervention developed by using linear programming: experience in Lombok, Indonesia. *Am J Clin Nutr* 2014; First published ahead of print December 24, 2014 as doi : 10.3945/ajcn.114.087775.

Hampel D, Shahab-Ferdows S, Domek JM, Siddiqua T, Raqib R, Allen LH. Competitive chemiluminescent enzyme immunoassay for vitamin B12 analysis in human milk. *Food Chem* 2014;153:60-5.

Jaiswal N, Melse-Boonstra A, Sharma SK, Srinivasan K, Zimmermann MB. The iodized salt programme in Bangalore, India provides adequate iodine intakes in pregnant women and more-than-adequate iodine intakes in their children. *Public Health Nutr* 2014;1-11 [Epub ahead of print].

Jaiswal N, Melse-Boonstra A, Thomas T, Basavaraj C, Sharma SK, Srinivasan K, Zimmermann MB. High prevalence of maternal hypothyroidism despite adequate iodine status in Indian pregnant women in the first trimester. *Thyroid* 2014;24:1419-29.

Jiang H, Li M, Wen LM, Hu Q, Yang D, He G, Baur LA, Dibley MJ, Qian X. Effect of short message service on infant feeding practice: findings from a community-based study in Shanghai, China. *JAMA Pediatr* 2014;168:471-8.

Khomsan A, Merita, Anwar F, Sukandar D, Riyadi H, Herawati T, Fatchiya A, Dewi M, Madanijah S, Syarif H. Impact of nutritional education on service quality in the integrated service posts (Posyandu). *Pakistan J Nutr* 2014;13:122-8.

Kulwa KB, Verstraeten R, Bouckaert KP, Mamiro PS, Kolsteren PW, Lachat C. Effectiveness of a nutrition education package in improving feeding practices, dietary adequacy and growth of infants and young children in rural Tanzania: rationale, design and methods of a cluster randomised trial. *BMC Public Health* 2014;14:1-16 doi: 10.1186/1471-2458-14-1077.

Macharia-Mutie CW, Omusundi AM, Mwai JM, Mwangi AM, Brouwer ID. Simulation of the effect of maize porridge fortified with grain amaranth or micronutrient powder containing NaFeEDTA on iron intake and status in Kenyan children. *Public Health Nutr* 2013;16:1605-13.

Mananga MJ, Kana-Sop MM, Nolla NP, Tetanye-Ekoe, Gouado I. Feeding practices and nutrition insecurity of infants and their mothers in Bangang rural community, Cameroon. *J Nutr Food Sci* 2014;4:264. doi: 10.4172/2155-9600.1000264.

Ndiku MH, Jara E, Sabaté J. Formative research on acceptability of pearl millet in rural eastern Kenya - a pilot study. *Sustainable Agriculture Research* 2014;3:1-8 doi:10.5539/sar.v3n4p1.

Nolla NP, Kana Sop MM, Mananga MJ, Ekoue T, Gouado I. Assessment of nutritional status of preschool children in the Bangang rural community, Cameroon. *Int J Biotechnol Food Sci* 2014;2:44-52.

Sunarti E. Empowering the care of family members to improve child nutritional status. *Int Federation for Home Economics* 2014;section 3:88-96.

Ukegbu PO, Anyika-Elekeh J, Uwaegbute AC, Ignatious J, Echendu CA, Nkwoala CC, Nzeagwu OC, Asumugha VU, Henry-Unaeze HN. Breastfeeding knowledge, beliefs and practices of mothers in Umuahia north local government: a baseline study. *Nigerian J Nutr Sciences* 2013;34:107-12.

Vossenaar M, Doak CM, Solomons NW. Challenges in the elaboration of a field interview instrument to capture information for the evaluation of adherence to the WHO/PAHO

Guiding Principles for Complementary Feeding of the Breastfed Child (Participation NF in the printing costs). Food and Nutrition Bulletin 2014;35:338-50.

Wijaya-Erhardt M, Muslimatum S, Erhardt JG. Effect of an educational intervention related to health and nutrition on pregnant women in the villages of central Java province, Indonesia. Health Education JI 2014;73:370-81.

2015

Ahmad SM, Hossain MB, Monirujjaman M, Islam S, Huda MN, Kabir Y, Raqib R, Lönnerdal BL. Maternal zinc supplementation improves hepatitis B antibody responses in infants but decreases plasma zinc level. Eur J Nutr (published online 25 July 2015) doi: 10.1007/s00394-015-0999-6.

Egbi G, Ayi I, Saalia FK, Zotor F, Adom T, Harrison E, Ahorlu CK, Steiner-Asiedu M. Impact of cowpea-based food containing fish meal served with vitamin C-rich drink on iron stores and hemoglobin concentrations in Ghanaian schoolchildren in a malaria endemic area. Food Nutr Bull 2015;36:264-275.

Helmyati S, Juffrie M, Rahayu ES, Istiti Kandarina BJ. A comparative study of gut microbiota profiles of children living in Kulon Progo and West Lombok. Pakistan J Nutr 2015;14:762-4.

Helmyati S, Rahayu ES, Kandarina BJ, Juffrie M. Sensory evaluation comparison of formulated fermented milk and commercial fermented milk: a new formulated fermented milk as food supplement for iron-deficiency anemia elimination program. Int J Curr Microbiol App Sci 2015;4:657-65.

Htet MK, Fahmida U, Thurnham DI, Hlaing LM, Akib A, Utomo B, Houghton LA. Folate and vitamin B12 status and dietary intake of anaemic adolescent schoolgirls in the delta region of Myanmar. Br J Nutr 2015;1-6 (doi:10.1017/S0007114515001609).

Jaikrishna N. Effect of iodine supplementation in Indian pregnant women on maternal and newborn thyroid function and cognitive development. Published Thesis Wageningen University, The Netherlands; 2015:241pp (ISBN 978-94-6257-333-8).

Kana Sop MM, Mananga MJ, Tetanye E, Goudao I. Risk factors of anemia among young children in rural Cameroon. Int J Curr Microbiol App Sci 2015;4:925-35.

Kana Sop MM, Namadi M, Tetanye E, Gouado I. Effect of Spirulina platensis supplementation on nutritional and biochemical parameters of under five years malnourished children from an orphanage in Douala, Cameroon. J Pharm Nutr Sci 2015;5:5-13.

Li M, Jiang H, Hu QZ, He GS, Wen LM, Dibley MJ, Baur L, Qian X. Text message to promote breastfeeding and obesity-protective eating behaviours in young children: 12 and 24 months BMI results. Obes Res Clin Pract 2014; 8 (Suppl. 1):58 (dx.doi.org/10.1016/j.orcp.2014.10.107).

Mangani C, Maleta K, Phuka J, Cheung YB, Thakwalakwa C, Dewey K, Manary M, Puumalainen T, Ashorn P. Effect of complementary feeding with lipid-based nutrient supplements and corn-soy blend on the incidence of stunting and linear growth among 6- to 18-month-old infants and children in rural Malawi. Matern Child Nutr 2015;11:132-43.

Nepal AK, Suwal R, Gautam S, Shah GS, Baral N, Andersson M, Zimmermann MB. Subclinical hypothyroidism and elevated thyroglobulin in infants with chronic excess iodine intake. Thyroid 2015;25:851-9.

Nguyen P, DiGirolamo A, Gonzalez-Casanova I, Pham H, Truong T, Nguyen S, Martorell

R, Ramakrishnan U. Breastfeeding practices are associated with child development in Vietnamese Children. The FASEB J 2015;29:1 (suppl 899.4) (abstract).

Rahmawati W, Muslimatun S, Wijaya-Erhardt M, Erhardt JG, Kurniawan A. Helminth infection and iron status of pregnant women in two geo-climatic areas in Indonesia. Malaysian J Nutr 2015;21:155-66.

Reinsma K, Bolima N, Fonteh F, Okwen P, Siapco G, Yota D, Montgomery S. Bobbi Be Best: the development and evaluation of an audio program and discussion guide to promote exclusive breastfeeding in Cameroon, Central Africa. Glob Health Promot 2015;1-13 (doi: 10.1177/1757975914556786) [Epub ahead of print].

Siddiqua TJ, Ahmad SM, Ahsan KB, Rashid M, Roy A, Rahman SM, Shahab-Ferdows S, Hampel D, Ahmed T, Allen LH, Raqib R. Vitamin B12 supplementation during pregnancy and postpartum improves B12 status of both mothers and infants but vaccine response in mothers only: a randomized clinical trial in Bangladesh. Eur J Nutr 2015 (published online February 2015) (doi: 10.1007/s00394-015-0845-x).

Sukandar D, Khomsan A, Anwar F, Riyadi H, Mudjajanto ES. Nutrition knowledge, attitude, and practice of mothers and children nutritional status improved after five months nutrition education intervention. Int J Sci: Basic Appl Res (IJSBAR) 2015;1-19.

Tafere G, Peterson S, Stoecker B. High variability of urinary iodine concentrations (UIC) and iodine in salt from rural households in Sidama zone, Southern Ethiopia. Eur J Nutr & Food Safety 2015;5:914-5 (abstract).

Ukegbu PO, Uwaegbute AC, Anyika JU, Echendu CA, Nkwoala CC. Assessment of lactation performance of mothers practicing exclusive breastfeeding in Abia State, Nigeria using test weighing technique. Pioneer Medical J 2015;5:16-24.

The publications are available free of charge upon request